



ISSN 2790 – 5985
eISSN 2790 – 5993

Agriculture College – Wasit University

Dijlah Journal of
Agricultural Sciences

Dijlah J. Agric. Sci., 4(3):56-66, 2025

Study effect of Al-Zubaidia Electric Station in soil pollution with heavy metals with different locations and wards in Wasit Province

Mahdi Wasmeiy Seheib Alaidi¹ Asmaa Hussein Allawe²

^{1,2}Department of Soil Sciences and Water Resources, College of Agriculture,
University of Wasit, Wasit, Iraq.

Corresponding author: malaiedy@uowasit.edu.iq

Abstract:

To study the pollutions with some heavy metals (Pb, Ni, Cr), a field study was applied to determine the effect of the Al-Zubaidia Electric Station in Wasit Province on soil pollutions by heavy metals in some area surroundings the industrials. Samples collected for three locations including 400, 900 and 1800 m away from pollutions sources, and the fourth site (control) was from the northwestern. Soil samples were collected at two depths (0-50) and (50-75) cm from southeastern, and northwestern areas. The result found an increase in total concentrations rate of heavy metals for lead, nickel and chromium in studied sites in the southeastern of the industrial at a depth of (50-0) cm, which is 400 m away from the asphalt industrial. These metals have exceeded the globally permissible limits, especially in the southern direction compared to the soil of the distant sites in the northwestern direction, especially at Depth (50-75) cm. Human activities were the main sources of the increase in pollutions standards values (CF and PLI). The CF values indicated the occurrence of large to very large pollution, especially in southeastern compared to northwestern areas, which recorded lower values. For PLI values, it indicated the occurrence of deterioration of the studied soils, especially in the southeastern area of the soils close to the laboratories. Therefore, the study recommends not to cultivate in the soils close to the Al-Zubaidia Electric Station and to use modern methods to reduce the gas emissions.

Keywords: Heavy metals, Al-Zubaidia Electric, Soil pollution.

Received:3/7/2024

Accepted:13/8/2025

Published:30/9/2024

دراسة تأثير محطة كهرباء الزبيديّة في تلوث التربة بالعناصر الثقيلة بأبعاد واتجاهات مختلفة في محافظة واسط

أسماء حسين علاوي²

مهدي وسمي صليب¹

^{1,2}جامعة واسط- كلية الزراعة- قسم علوم التربة والموارد المائية

الخلاصة

لدراسة تأثير تلوث التربة ببعض العناصر الثقيلة المنبعثة من محطة كهرباء الزبيدية الحرارية في محافظة واسط (الرصاص، النيكل، الكروم). اجريت دراسة ميدانية لتقدير العناصر الثقيلة في المناطق المحيطة بالمعامل، إذ جمعت نماذج تربة من ثلاثة مواقع قريبة من المحطة وعلى بعد 400 و 900 و 1800 م من مصادر التلوث فضلا عن اخذ نماذج تربة على بعد 3500 م ليمثل معاملة المقارنة، وبعمقين (0-50 سم و 50-75 سم) ومن اتجاهين مختلفين (جنوبي شرقي وشمالي غربي). اظهرت نتائج الدراسة ارتفاع التراكم الكلي للعناصر الثقيلة عن الحدود المسموح بها عالميا لكل من الرصاص والنيكل والكروم في مواقع الدراسة لاسيما الجزء الجنوبي الشرقي عند العمق 0-50 سم وعلى بعد 400 م قياسا بالجزء الشمالي الغربي عند نفس البعد والعمق، بينما انخفضت تراكيز العناصر الثقيلة كلما ابتعدنا عن مصدر التلوث والتمثلة بالابعاد 900 و 1800 م عند العمق 50-75 م ولكلا الاتجاهين. وبينت الدراسة ايضا الى ارتفاع في مؤشرات التلوث والتمثلة بمؤشر عامل التلوث (CF) ومؤشر حمل التلوث (PLI)، إذ كان التلوث شديد الى شديد جدا عند الاتجاه الجنوبي الشرقي لاسيما الموقع 400 م بعمق 0-50 سم قياسا بالاتجاه الشمالي الغربي عند نفس الموقع والعمق، بينما كانت مؤشرات التلوث اقل عند الابعاد 900 و 1800 م ولكلا الاتجاهين من محطة الكهرباء الحرارية.

الكلمات المفتاحية: العناصر الثقيلة، كهرباء الزبيدية، تلوث التربة.

1. Introduction

Environmental pollution is any quantitative or qualitative change in the chemical, physical and biological properties of one or more metals of the environment such as air, water and soil, which leads to significant harm that affects human health and all living organisms. The harm of pollution results from different human activities, most notably industrial activities [Lechukwu *et al*:2021]. Everything that pollutes the air and water in turn affects the soil and contributes to its pollution, as water and air fill the largest part of the interstitial spaces in the soil [Radomirovic *et al*:2020].

The problem of pollutions is considered one of important problem that began to takes on the serious environmental dimension, especially the industrial revolutions that occurred in Europe and the industrial expansion that occurred, which was accompanied by the emergence of serious pollution in the ecosystems. The problem of pollutions was not a new problem for the ecosystems, but the increase in the severity of pollution is new in our time, and the risk of environmental pollution is no less dangerous than the occurrence of wars and conflicts [Moghtaderi *et al*:2018].The problem of pollution, especially industrial pollution, is characterized by many characteristics that cause deterioration in the quality of the environment surrounding industrial complexes, oil refineries, power plants and electrical stations.

These characteristics are the non-compliance of their location and the ages of these facilities, and their lack of mean to treat pollutions, in addition to their lack of environmentally appropriate technology. All of this leads to a significant increase in the concentrations of pollutants resulting from them [Spreadbury *et al* :2021].Therefore, this study aimed to estimate the effect of emissions from asphalt industrial on the accumulations of the heavy metal in soil near the industrial, in addition to evaluating the levels of soil pollutions with heavy metal (Pb, Co and Cd) resulting from the combustion of fuel used in Al-Zubaidia Electric Station in two different directions and depths and different locations and comparing them with global determinants.

2. Materials and Methods

2.1. Study Area

The samples were collected is located in Al-Zubaidia District, Wasit Province, which includes many Al-Zubaidia Electric Station and the area is surrounded by agricultural lands. The area divided to two areas, the southeast and the northwestern. Soil samples collected on 25/6/2024 from southeastern, and northwestern areas at three distances from the pollution source. The dimensions were 400 - 900 - 1800 m, respectively, for the southeastern part and the northwestern part. The comparison sample was taken at a distance of 3500 m and at two depths for each site, the first (0-50) cm and the second (50-75) cm, with three sample for each site.

2.2. Estimations of Heavy Metals in the Soil

Heavy metals (Cr - Pb - Ni) were estimated using a mixture of perchloric and sulfuric acids using an absorption device Atomic Absorption Spectrophotometer (AAS) as described in the method [Wu *et al*:2022].

2.3. Calculating Environmental Pollution Indicators for Soil

Contamination factor (CF): [Krishna and Mohan:2014]

$$Cf = c_m \text{ sample} / c_m \text{ background}$$

Cf = cm samples. cm backgrounds

Cf = Contaminations factors

Cm = Total concentrations of heavy metals in soil samples (mg/kg⁻¹ soil)

Cm backgrounds = Total concentrations of heavy metals in control soil samples (mg/kg⁻¹).

2.4. Pollution Load Index (PLI)

$$PLI = (CF_1 \times CF_2 \times CF_3 \dots CF_n)^{1/n}$$

Cf = Contamination factors for the first, second, third metals... etc.,

n = Number of heavy metal study.

3. Results and Discussion

3.1. Total Concentrations of Heavy Metals in the Study Soils

3.1. Total Lead Pb⁺²

Table (1) indicates Total lead concentrations in soil for the three studied sites (400, 900 and 1800) m and for both directions (southeast and northwest) and for the two dimensions (0-50) cm and (50-75) cm. The values for the southeast part reached (307.33 and 261.55) mg Pb/kg⁻¹ for the dimensions 400 and 900 m respectively compared to third dimensions (1800) m and the comparison samples (3500) m, that reached values of 114.85 and 42.79 mg Pb/kg⁻¹ respectively. The result in Table (1) showed that values achieved for the totals leads concentrations in soil at the first depths (0-50) cm were higher compared to the second depth (50-75) cm for the same direction and dimensions, as the values reached (195.64 and 167.62) mg Pb/kg⁻¹ for the first, and second depths, respectively. As for the overlap between the dimension and depth, the high value achieved at the dimension (400) m for the depth (0-50) cm in the south-eastern direction, which was (321.81) mg Pb/kg⁻¹, while low value was achieved at the dimension (1800) m for the depth (50-75) cm, which was (97.96) mg Pb/kg⁻¹. The result of statistical indicated the existence of significant differences for each of the dimension, depth and overlap between them. The result recorded that the value of lead concentrations in the soil in the northwest direction and for all dimension, and depth studied were lower compared to southeast direction.

The average values reached (109.48, 93.91 and 70.05) mg Pb/kg⁻¹ for dimension (400, 900 and 1800) m, respectively, compared to the control reached (42.79) mg Pb/kg⁻¹ soil while the first depth (0-50) cm achieved higher values compared to second depth (50-75) cm. The result of statistical indicated the presence of significant differences for each of dimension, depth, and the

interactions between them. An increase in concentrations of lead in soil at site was close to asphalt industrials, and in both the southeast and northwest directions, according to the permissible global determinant (WHO, 2007) ($100 \text{ mg Pb/kg}^{-1}$ soil).

This may be because to several reasons, including that sites close to asphalt industrial are affected by the heavy metals and gases released by these industrial into the atmosphere, and these gases are deposited in sites close to the industrial, which effectively contributes to the pollution of the surrounding soil by increasing the concentrations of heavy metal in it. In addition, the level of lead in surface layer of the soil is affected by the rate of accumulation and deposits from the atmosphere emitted from the industrial, and this accumulation gradually decreases the further we move away from the industrial site, in addition to the role of the prevailing winds in the region (north-west), which contribute to the transfer of pollutants from the source to nearby sites and their depositions on the soil surfaces. The result was agreed with [Kreshna and Mohan:2014, Saeedi and Salmanzadeh:2012, Kong *et al*:2011, Tran *et al*,2022], as their studies indicated that the concentrations of lead in soil increases in site close to pollutions sources.

Table 1. Concentration of lead Pb (mg Pb/kg^{-1} soil) in the studied sites.

Site	Distance (m)	Depth (cm)		Average
		0-50	50-75	
southeast	400	321.81	292.86	307.33
	900	284.81	238.29	261.55
	1800	131.75	97.96	114.85
Control	3500	44.21	41.37	42.79
Average		195.64	167.62	---
LSD (0.05)		Distance	Depth	Distance* Depth
		*2.27	*2.08	* 3.93
Northwest	400	119.14	99.83	109.48
	900	101.16	86.67	93.91
	1800	76.12	63.99	70.05
Control	3500	44.21	41.37	42.79
Average		85.16	72.96	---
LSD (0.05)		Distance	Depth	Distance* Depth
		* 1.79	* 1.46	* 2.30
Indicators	WHO , 2007		$100 \text{ mg /Pb kg}^{-1}$	
	Kabata-Pendias , 2011		$300 \text{ mg /Pb kg}^{-1}$	

3.2. Total Nickel Concentration Ni in the Study Soils

The results and statistical analysis indicates that there was variations in the total nickel concentrations in soil from one site to another as a result of proximity and distance from the pollution source and the overlap between them. The result showed that distances from pollutions sources affected the soil content of the total nickel metals. There were significant differences in southeastern part of laboratories, as first site has recorded the highest value of the metals concentration of ($100.63 \text{ mg Ni/kg}^{-1}$ soil) compared to control treatment, which achieved the lowest value ($9.56 \text{ mg Ni/ kg}^{-1}$ soil) (Table 2).

This is agreed with [Kabir *et al*,2022] that the concentration of the total nickel metals increases in sites close to brick industrial. He attributed the reason for this to the fact that the gases emitted from brick industrial settle in sites close to the industrial, which increases the concentrations of the total metals in the soil. Also, [Zheng *et al*,2020] found that soils near thermal stations and adjacent to refineries in Baiji. The result showed that concentrations of total nickel in the soil is affected by distance as well as wind direction, and the concentration of the metals decreases the further away we are from the sources of pollutions.

The results in Table (2) showed significant difference in depth in concentrations of total nickel in studied soil sample. The first depth (0-50) cm recorded the highest value of nickel, amounting to (83.35) mg Ni/ kg⁻¹, with increase rate of 40% compared to second depth (50-75) cm, which recorded the lowest value, amounting to (59.62) mg Ni/kg⁻¹. These results are consistent with [Al-Saad et al,2018], concluded that the concentration of the metals is higher in the first depth compared to concentrations of metals in the second depth. The result of two-way interactions between dimensions, and depths had significant effects on the concentrations of total nickel in the soil, as the first depth (0-50) cm with the first dimension (400) m recorded the highest value of the metals concentration, reaching (117.14) mg Ni/kg⁻¹, compared to the control samples (3500) m, and the second depth (50-75) cm, where it reached (8.46) mg Ni/kg⁻¹.

Table 2. Nickel concentrations Ni (mg Ni/kg⁻¹) in the soil.

Site	Distance (m)	Depth (cm)		Average
		0-50	50-75	
southeast	400	117.14	84.12	100.63
	900	109.8	80.94	95.37
	1800	95.78	64.95	80.36
Control	3000	10.67	8.46	9.56
Average		83.35	59.62	---
LSD (0.05)		Distance	Depth	Distance* Depth
		* 1.78	* 1.29	* 2.08
Northwest	400	80.66	54.96	67.81
	900	74.82	49.28	62.05
	1800	56.64	37.6	47.12
Control	3500	10.67	8.46	9.56
Average		55.69	37.57	---
LSD (0.05)		Distance	Depth	Distance* Depth
		* 1.47	* 1.06	* 1.89
Indicators	WHO , 2007		50 mg Ni/ kg ⁻¹	
	Kabata-Pendias , 2011		60 mg Ni/ kg ⁻¹	

For the results of the northwestern part of the Al-Zubaidia Electric Station, the results of Table (2) showed differences in effect of distance, and proximity as well as the overlap between depth and distance in the concentrations of the nickel metals in the soil. It notes that the nickel metals recorded the highest value in the first sites close to the pollution source and the first depth (0-50) cm, as the value reached (80.66) mg Ni/kg⁻¹ soil, compared to second depth (50-75) cm and the control samples, where the value reached (8.46) mg Ni/kg⁻¹ soil. When comparing these results with the permissible global determinants (WHO, 2007), our result find that concentrations of nickel metals in the southeastern and northwestern parts exceeded the permissible limits.

3.3. Total Chromium Concentrations(Cr) in the Study Soil

The result showed the effect of distances, and proximity to Al-Zubaidia Electric Station. The total chromium concentration recorded its highest value within the first dimension, reaching (88.99) mg Cr/kg⁻¹, compared to control treatment, that recorded low value (4.48) mg Cr/kg⁻¹ (Table 3). The reason for this may be attributed to the gases, and vapours rising from the Al-Zubaidia Electric Station and heavy metals they carry, which contribute to increasing the concentration of heavy metals in areas close to pollutions sources.

The results agreed with [Bedeh *et al*,2022 ,Davies,1992] that the concentration of chromium has increased in the Dora area and attributed reason to activity of the Dora refinery and power station, which contributes to increasing the amount of pollution. The results in Table (3) showed significant effects in depth in total concentrations of total chromium of soil sample within the

southeastern part of the Al-Zubaidia Electric Station. The first depth (0-50) cm gave the highest value of chromium, reaching 54.12 mg Cr/kg⁻¹ soil, compared to second depth (50-75) cm, which recorded a value of 40.84 mg Cr/kg⁻¹. The reason for the decrease may be attributed to the accumulation of heavy metals in surface layer of the soil, in addition, to the lack of rainfall, which works to wash away heavy metals and transfer them to the lower soil horizons.

Table 3. Chromium (Cr) concentration (mg Cr/kg⁻¹ soil).

Site	Distance (m)	Depth (cm)		Average
		0-50	50-75	
southeast	400	101.16	76.83	88.99
	900	79.21	61.91	70.56
	1800	31.18	20.62	25.90
Control	3500	4.93	4.03	4.48
Average		54.12	40.84	---
LSD (0.05)		Distance	Depth	Distance* Depth
		* 2.50	* 1.63	* 3.42
Northwest	400	34.54	26.59	30.56
	900	22.78	11.91	17.34
	1800	12.74	6.94	9.84
Control	3500	4.93	4.03	4.48
Average		18.75	12.36	---
LSD (0.05)		Distance	Depth	Distance* Depth
		* 1.37	* 1.01	* 1.83
Indicator	Kabata-Pendias , 2011		200 mg Cr/ kg ⁻¹	

For the interactions for distance, and depth, it gave significant effects on the concentration of the total chromium metals in soil, as the first depth (0-50) cm with the first dimension gave highest values for chromium metals (101.16) mg Cr/kg⁻¹ soil compared to second depths (50-75) cm with the control, which gave 4.03 mg Cr/kg⁻¹, with a clear significant decrease from the rest of the values. The results in Table (3) showed significant differences in the northwestern part of the dimension and depth and their bilateral interactions in the concentrations of total chromium metals in the soil. The chromium metals in the first site recorded a value of (30.56) mg Cr/kg⁻¹, compared to control, that was (4.48) mg Cr/ kg⁻¹ soil. The results agreed with Tran et al. (2022). It was found that the effect of depths in taking samples had significant effects on the concentrations of the chromium metals. The first depth (0-50) cm gave the highest values was (18.75) mg Cr/kg⁻¹, with an increase rate of 51% compared to the second depth (50-75) cm, that recorded (12.36) mg Cr/kg⁻¹ soil.

The interactions between distance, and depth had significant effects on the concentrations of total chromium, as the first depth (0-50) cm with the first dimension (400) m has recorded high values of (34.54) mg Cr/kg⁻¹ soil compared to second depth (35-70) cm and the control recorded the lowest value of (4.03) mg Cr/kg⁻¹. Through the results of our study, the results find that the concentrations of total chromium in soil increased in southeastern area compared to northwestern parts. This is due to nature of the prevailing wind in the region (north-west), and when comparing these result with the permissible limit according to [Gharaybeh, 2010, Hakanson, 1980], it find that the concentrations of total chromium was below permissible limit in all studied sites and for both the southeastern and northwestern parts.

3.4. Pollution Indicator for Heavy Metals in the Study Soil

Pollutions Factor

Table 4. Pollutions factor value.

Depth (cm)	Site	Distance (m)	Pb	Ni	Cr
0-50	southeast	400	7.28	10.98	2052
		900	6.44	10.29	16.07
		1800	2.98	8.98	6.32
	Northwest	400	2.69	7.56	7.01
		900	2.29	7.01	4.63
		1800	1.72	5.31	2.58
50-75	southeast	400	7.08	9.94	19.06
		900	5.76	9.57	15.36
		1800	2.37	7.68	5.12
	Northwest	400	2.41	6.49	6.59
		900	2.09	5.82	2.95
		1800	1.55	4.44	1.72

3.5. Pollution Factor (CF) for Lead Metal Pb

The results showed the value of the pollutions factors for lead metal of sites included in the southeastern, and northwestern parts of the asphalt industrial. In the southeastern part, the first site (400) m and for the first depth (0-50) cm recorded the high value of the pollutions factors, which amounted (7.28), compared to third sites (1800) m and for the same first depth, which recorded the lowest value, which amounted to (2.98) (Table 4). The pollutions factors in first sites indicates the occurrence of very severe pollutions of lead metal, while the value of pollutions factors in third sites indicates the occurrence of medium pollutions according to classifications[19]. However, when comparing the depths, we find that the pollution factor has decreased significantly at the second depth (50-75) cm compared to the first depth (0-50) cm. For the northwestern part of the asphalt industrial, the result showed a decreases in values of the pollutions index compared to southeastern in Table (4). It is found that value of the pollutions factor index for the first, the second, and the third dimension were (2.69, 2.29 and 1.72) respectively, and this indicate the occurrence of moderate pollution with lead according to the classification [Kabata,2011].

The results of the study agreed with [20] that an increases in value of the pollutions factors for lead in sites close to the Dora refinery. Tran *et al.*, (2022) indicates that reason to impact of the brick plant on soil pollutions. The result showed an increase in the value of the pollutions factors in the south eastern compared to the northwestern of Al-Zubaidia Electric Station, and this is attributed to the natures of prevailing wind (northwest) in regions, which work to carry pollutant resulting from industrials activities to southeastern arae of the asphalt industrial.

3.6. Pollution Factor (CF) for Nickel (Ni)

The results of Table(4) showed that there is a variation in level of the pollutions factor for the nickel according to the proximity, and distances from the pollution source, as well as according to the location, and depth of taking soil sample in the southeastern and northwestern parts of the asphalt industrial. In the southeastern of the asphalt industrials, the high values of the pollutions factors was recorded in the first site (400) m and the first depth (0-50) cm, where it reached (10.98), compared to the third site (1800) m and the second depth (50-75) cm, which recorded the lowest value of (7.68). It notes that the values of the pollutions factors in the two sites indicate the occurrence of very severe pollutions, according to the classifications[Miskowiec *et al*,2015]. In the northwestern part, the pollutions factors value decreased compared to the southeasterns part, as the first site (400) m and first depth (0-50) cm recorded (7.56), while the pollutions factors value decreased in third site (1800) m and the second depth (50-75) cm, which gave the

lowest value of (4.44). The results indicate that there were very severe pollutions in the first site and severe pollutions in third site with nickel in the northwestern part of the asphalt industrial, according to the classifications [Saeedi and Salmanzadeh, 2012]. Through the results find that the pollutions factor values have increased in sites close to the pollutions sources, and this is attributed to the activity of asphalt industrial in the region and the accompanying accumulations of heavy metal in sites close to the pollutions source as a result of gases and vapours emitted from asphalt industrial.

These results are consistent with [Tomlinson *et al*, 2015], which found that values of the pollutions factors for the nickel metal increases in places close to the pollution source and with different types of pollution sources. In general, we find that the value of the pollutions factors have increased at the first depth (0-50) cm compared to second depth (50-75) cm for all dimensions, and both directions.

3.7. Pollution Factor (CF) for Chromium Metal (Cr)

The result showed that there is a variation in the value of the pollutions factors for chromium metals in soil of studied sites. In the southeastern of the asphalt industrial and in first site (400) m, the high values of the pollutions factors were recorded (20.52), and the lowest value was recorded in the third site (1800) m, where it reached (6.32) and for the first depth (0-50) cm Table (4).

Through the values of the pollutions factors, soil of sites is classified, according to standards [WHO, 2007], as very highly polluted. The reason for this is attributed to the high concentrations of total chromium in sites near asphalt industrial as a result of gaseous emissions containing heavy metal that are transported by air, and fall on the soil near the source of pollutions. This is consistent with what was indicated by [Kong *et al*, 2011] about high value of the pollutions factors for heavy metal in the soil of sites near industrial facilities.

In the northwestern of the asphalt, the pollutions factors found low value compared to southeastern part, the first site was (7.01) and the third site recorded (2.58) and for the first depth (0-50) cm, according to the classifications (Tomlinson *et al.*, 1980), the soil was classified from moderately polluted to very severely polluted soil. We note that the result of found that the values of the pollutions factors recorded a decrease at the second depth (50-75) cm compared to the first depth (0-50) cm for both directions. It notes an increase in the values of the pollutions factors in the southeastern compared to northwestern. The reason for this is attributed to the nature of the prevailing north-westerly wind in the regions, which work to carry pollutant from pollutions sources to the southeastern, which lead to an increase in pollutants in it compared to located opposite to the directions of the prevailing winds.

3.7. Pollution Load Index (PLI)

The result in Table (5) indicated variations in the value of the pollutions Load Index (PLI) for studied heavy metal of lead, nickel, cadmium, cobalt, and chromium. The southeastern part of the asphalt plant found that the high value of Pollution Load Index was at the first site (400) m and for the first depth (0-50) cm, which amounted to (11.85) compared to the third site (1800) m and the second depth (50-75) cm in the northwest direction, which recorded the low values of the Pollution Load Index, which was (2.39). From the result, the average value of the Pollution Load Index in the southeastern, and northwestern parts of the studied soil were greater than 1, and this indicates the state of deterioration of the studied soils. The reason for this is attributed to the gases, and the vapors emitted from the asphalt industrial as a results of their work, which contains large amounts of heavy metal that contribute to the deterioration of the soil surrounding those industrial.

Table 5. Pollution Load Index (PLI) value for the studied metals.

Depth (cm)	Distance from pollution source (m)	Southeast	Northwest
0-50	400	11.85	5.67
	900	9.88	4.44
	1800	5.96	3.07
50-75	400	10.63	4.64
	900	8.54	3.50
	1800	4.72	3.39
Pollution Load Index	1>PLI Site is unpolluted	PLI=1 Site is on the verge of deterioration	PLI >1 quality deterioration

The results agreed with [AL-Saad and Dana,2018], as they found an increases in the value of the pollutions load index in the soil of urban sites, and attributed reason for this to different human activity as well as power station, electric power generator and oil fields and the untreated pollutants they release into the soil that contribute to increasing the amount of heavy metals in the soil and its deterioration. The study also agreed with what was found by [Duong and Lee,2011] in a study of the soil of sites near industrial facilities, as the results of the pollutions load index were greater than 1 and attributed the reason for this to the gases, and the vapours emitted from power stations that contribute to increasing the percentage of heavy metal in the studied sites.

4. Conclusions

The study concludes that the CF values indicated the occurrence of large to very large pollution, especially in southeastern compared to the northwestern parts, which recorded lower values. For PLI values, it indicated the occurrence of deterioration of the studied soil, especially in the southeastern of soils close to the laboratories.

References

- [1] Ilechukwu, I., Osuji, L.C., Okoli, C.P., Onyema, M.O. and Ndukwe, G.I., 2021. Assessment of heavy metal pollution in soils and health risk consequences of human exposure within the vicinity of hot mix asphalt plants in Rivers State, Nigeria. *Environmental monitoring and assessment*, 193(8), p.461.
- [2] Radomirović, M., Ćirović, Ž., Maksin, D., Bakić, T., Lukić, J., Stanković, S. and Onjia, A., 2020. Ecological risk assessment of heavy metals in the soil at a former painting industry facility. *Frontiers in Environmental Science*, 8, p.560415.
- [3] Moghtaderi, T., Mahmoudi, S., Shakeri, A. and Masihabadi, M.H., 2018. Heavy metals contamination and human health risk assessment in soils of an industrial area, Bandar Abbas–South Central Iran. *Human and Ecological Risk Assessment: An International Journal*, 24(4), pp.1058-1073.
- [4] Spreadbury, C.J., Clavier, K.A., Lin, A.M. and Townsend, T.G., 2021. A critical analysis of leaching and environmental risk assessment for reclaimed asphalt pavement management. *Science of the Total Environment*, 775, p.145741.
- [5] Page ,A.L.;R.H.Miller and D.R Keeney (1982) . *Methods of soil analysis, part 2* 2 nd ASA Inc.Madison, Wisconsin.1158 pp.
- [6] Mirzaei, M., Baalousha, M., Mousavi, R., Mirzaei Aminiyan, F., Hosseini, H. and Heydariyan, A., 2018. The ecological risk, source identification, and pollution assessment of heavy metals in road dust: a case study in Rafsanjan, SE Iran. *Environmental Science and Pollution Research*, 25, pp.13382-13395.
- [7] Wu, G., Wang, L., Yang, R., Hou, W., Zhang, S., Guo, X. and Zhao, W., 2022. Pollution characteristics and risk assessment of heavy metals in the soil of a construction waste landfill site. *Ecological Informatics*, 70, p.101700.

- [8] Krishna, A.K. and Mohan, K.R., 2014. Risk assessment of heavy metals and their source distribution in waters of a contaminated industrial site. *Environmental Science and Pollution Research*, 21(5), pp.3653-3669.
- [9] Saeedi, M., Li, L.Y. and Salmanzadeh, M., 2012. Heavy metals and polycyclic aromatic hydrocarbons: pollution and ecological risk assessment in street dust of Tehran. *Journal of hazardous materials*, 227, pp.9-17.
- [10] Kong, S., Lu, B., Ji, Y., Zhao, X., Chen, L., Li, Z., Han, B. and Bai, Z., 2011. Levels, risk assessment and sources of PM10 fraction heavy metals in four types dust from a coal-based city. *Microchemical Journal*, 98(2), pp.280-290.
- [11] Tran, T.S., Dinh, V.C., Nguyen, T.A.H. and Kim, K.W., 2022. Soil contamination and health risk assessment from heavy metals exposure near mining area in BacKan province, Vietnam. *Environmental geochemistry and health*, 44(4), pp.1189-1202.
- [12] Kabir, M.H., Kormoker, T., Shammi, R.S., Tusher, T.R., Islam, M.S., Khan, R., Omor, M.Z.U., Sarker, M.E., Yeasmin, M. and Idris, A.M., 2022. A comprehensive assessment of heavy metal contamination in road dusts along a hectic national highway of Bangladesh: Spatial distribution, sources of contamination, ecological and human health risks. *Toxin Reviews*, 41(3), pp.860-879.
- [13] Zheng, N., Hou, S., Wang, S., Sun, S., An, Q., Li, P. and Li, X., 2020. Health risk assessment of heavy metals in street dust around a zinc smelting plant in China based on bioavailability and bioaccessibility. *Ecotoxicology and Environmental Safety*, 197, p.110617.
- [14] Al-Saad, Hamid T and DuhaS.Karem.(2018).Envvironmental Assessment of Heavy Metals in Soil of West Quarna-2 Oil field at Basrah-Southern Iraq.J.of Pet. Res.and Stu.,No.19.
- [15] Bedeeh , Riyadh, S. and Salwa ,J. Fakher(2022). Evaluation of Soil Pollution With Some Heavy Metals for the Area Around the Al-Ghaf Oil Field. *International J.of Agri.andstatisical Sci.* Vol.18 Supplement 1,2022.
- [16] Davies,B.E.,(1992).Inter-relationships between soil properties and the uptake of cadmium,copper,lead and zinc from contaminated soils by radish (*Raphanussativus L.*). *Water,air,and soil pollution* , 63(3-4):331-342.
- [17] Gharaybeh.K.M.(2010).Environmental pollution-concept ,forms and how to reduce the severity of it .*Journal of Environmental Studies*. Volume 3:121-133.
- [18] Hakanson L (1980). An Ecological risk index for aquatic pollution control.A sediment-logical approach. *Water Res* .14:975-1001.Doi :10.116/043-1354(80)90143-8.
- [19] Jackson , M. L. 1958 . *Soil chemical analysis* . N. J. Englewood Cliffs :Prentice-Hall Inc.
- [20] Kabata-Pendias ,A (2011).Trace elements in s soils and plants, Fourth – Edition . by Taylor and Francis Group,LLC.ISBN 978-1-4200-9368-1.
- [21] Miskowiec .P;A, Laptas , and K, Zieba.(2015).Soil poilution with heavy metais in industrial and agricultural areas: a case study of Olkusz District . *Journal of Elementology* ISSN 1644-2296.
- [22] Tomlinson, D.C.; J.G. Wilson; C.R.Harris and D.W.Jeffrey.(1980). Problems in the assessment of heavy metals in estuaries and the formation pollution index.Helgol. Mar.Res.33,566-575.
- [23] WHO(World Health Organization).(2007). Health risks of heavy metals from long range Trans boundary air pollution. <http://www.euro.who.int/pubrequest> .
- [24] Duong, T.T. and Lee, B.K., 2011. Determining contamination level of heavy metals in road dust from busy traffic areas with different characteristics. *Journal of environmental management*, 92(3), pp.554-562.